



# **CONSTIPATION**

## **CONSTIPATION & ASSOCIATED SYMPTOMS**



Unit Size: H 10.5 cm x W 5 cm x D 5 cm Weight 56 g UPC 6 65808 11007 0

#### **COMPOSITION:**

Medicinal ingredients	Properties of each ingredient
Alumina 6CH	Atonic constipation, slow digestive transit.
Causticum 6CH	Digestive transit paralysis with constipation, defecation difficulties, loss of rectal sensitivity.
Lycopodium 6CH	Infant digestive problems, gas emission, meteorism in umbilical region, thin and elongated stools.
Magnesia muriatica 6CH	Hepatic and digestive syndrome with constipation, dehydrated stool. Milk intolerance.
Nux vomica 6CH	Constipation with ineffectual urging and tenesmus. The stools are hard and incompletely evacuated. Abdomen sensitivity during walking. Painful and difficult digestion.
Sepia officinalis 6CH	Sensation of abdominal emptiness. Atonic constipation. Bearing down in the abdomen.
Bryonia alba 6CH	The abdominal wall is tense, sensitive to touch and calmed by pressure. Painful bloating, constipation with large, hard and dry stools.

## Non medicinal ingredients:

Purified water, sorbitol, glycerin, sodium benzoate, potassium sorbate, citric acid.

#### Indications:

KIDZ CONSTIPATION oral solution is recommended for the treatment of constipation and associated symptoms: constipation due to dehydrated stools, small and insufficient, with difficult and prolonged evacuation. Sensitive and tender abdomen.

**Dosage:** To be taken orally

Infants 0-11 months old: 1 to 5 drops, 1 to 3 times per day. Children 1-5 years old: 5 to 10 drops, 1 to 3 times per day. Children 6-11 years old: 5 to 15 drops, 1 to 3 times per day. At the start of the treatment, take every 30 minutes, up to 10 times per day,

then return to general dosage. To be taken directly under the tongue with the dropper or to melt with some water before drinking.

## **Duration of use:**

3 to 4 days.

## Warnings:

If symptoms persist more than 4 days or are accompanied by fever, consult a healthcare practitioner.

## Incompatibility with other medicine:

No known incompatibility with other medicines.

